

University of Dundee

## Reduction in post-operative acute kidney injury following a change in antibiotic prophylaxis policy for orthopaedic surgery

Walker, Heather; Patton, Andrea; Bayne, Gwen; Marwick, Charis; Sneddon, Jacqueline; Davey, Peter

*Published in:*  
Journal of Antimicrobial Chemotherapy

*DOI:*  
[10.1093/jac/dkw166](https://doi.org/10.1093/jac/dkw166)

*Publication date:*  
2016

*Document Version*  
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

### *Citation for published version (APA):*

Walker, H., Patton, A., Bayne, G., Marwick, C., Sneddon, J., Davey, P., Nathwani, D., & Bell, S. (2016). Reduction in post-operative acute kidney injury following a change in antibiotic prophylaxis policy for orthopaedic surgery: an observational study. *Journal of Antimicrobial Chemotherapy*, 71(9), 2598-2605.  
<https://doi.org/10.1093/jac/dkw166>

### **General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

[Type text]

**Reduction in postoperative Acute Kidney Injury following a change in antibiotic prophylaxis policy for orthopaedic surgery – an observational study**

Heather Walker<sup>1</sup>, Andrea Patton<sup>2</sup>, Gwen Bayne<sup>3</sup>, Charis Marwick<sup>2</sup>, Jacqueline Sneddon<sup>3</sup>, Peter Davey<sup>2</sup>, Dilip Nathwani<sup>3</sup>, Samira Bell<sup>1</sup>

<sup>1</sup>Renal Unit, Ninewells Hospital, Dundee, DD1 9SY, Scotland, <sup>2</sup>Population Health Sciences, School of Medicine, University of Dundee, Dundee, DD2 4BF, Scotland, <sup>3</sup>Scottish Antimicrobial Prescribing Group, Scottish Medicines Consortium, Glasgow, G1 2NP, Scotland.

**Correspondence:**

**S Bell**

**Tel: 00 44 1382 633913**

**Fax: 00 44 1382 632327**

**E-mail: [samira.bell@nhs.net](mailto:samira.bell@nhs.net)**

**Short running title:** Reduced AKI in orthopaedic surgery

## Abstract

Objectives: Evidence has shown that a prophylactic antibiotic regimen of flucloxacillin and gentamicin for orthopaedic surgery was associated with increased rates of post-operative acute kidney injury (AKI). This resulted in changes in national antibiotic policy recommendation for orthopaedic surgical prophylaxis. This study aimed to assess whether this change from flucloxacillin and gentamicin to co-amoxiclav was associated with changes in the rates AKI and *Clostridium difficile* infection (CDI).

Methods: An observational study and interrupted time series analyses were used to assess rates of post-operative AKI separately in patients undergoing neck of femur (NOF) repair and other orthopaedic operations which required antibiotic prophylaxis. Incidence rate ratios were used to evaluate changes in CDI rates.

Results: Following the change in policy, from flucloxacillin and gentamicin to co-amoxiclav, there was a relative change in rates of post-operative AKI of -63% (95% CI -77% to -49%) at 18 months in the other orthopaedic surgeries group. In the NOF repair group, there was no change in the rate of post-operative AKI -10%, (95% CI -35% to 15%) at 18 months. The incident rate ratio for CDI in other orthopaedic operations was 0.29 (95%CI 0.09 to 0.96) and for NOF repair patients was 0.76 (95%CI 0.28 to 2.08).

Conclusions: The use of co-amoxiclav for antibiotic prophylaxis in orthopaedic surgery was associated with a decreased rate of post-operative AKI compared to flucloxacillin and gentamicin and was not associated with increased rates of CDI.

## Introduction

Surgical site infections (SSI) are a preventable healthcare associated infection. They are associated with a significant increase in morbidity, mortality, length of hospital stay and both socioeconomic and health related costs.<sup>1, 2</sup> Prophylactic antibiotics have been shown to reduce the rate of SSI in patients undergoing joint replacement surgeries<sup>3</sup> and are recommended for orthopaedic implant surgeries.<sup>4</sup> However, their use can also have associated adverse effects and unintended consequences, such as allergy,<sup>5</sup> the development of antibiotic resistance among organisms,<sup>6</sup> acute kidney injury (AKI)<sup>7-10</sup> and *Clostridium difficile* infection (CDI).<sup>11, 12</sup>

AKI is reported to be an independent risk factor for mortality, as well as being a marker of co-existing pathologies,<sup>13-16</sup> and is a predictor of poor prognosis in hospitalised patients.<sup>17-19</sup> AKI is associated with increased risk of the development and progression of chronic kidney disease (CKD), increased risk of readmission,<sup>20</sup> longer hospital stays and higher long term mortality rates,<sup>19</sup> even in patients with transient changes in renal function.<sup>20</sup> AKI therefore has associated important long term socioeconomic and public health effects.

CDI is a healthcare-associated infection with significant associated morbidity and mortality<sup>21</sup> that has been under mandatory surveillance in Scotland since 2006,<sup>22</sup> and in England since 2004.<sup>23</sup> Antibiotic use is associated with an increased risk of CDI for at least 3 months following administration.<sup>24</sup> In order to restrict the use of antibiotics associated with high risk of CDI<sup>25</sup> NHS Tayside orthopaedic antibiotic prophylaxis was changed from cefuroxime to gentamicin and flucloxacillin in October 2008. Following this, several studies indicated that prophylaxis with flucloxacillin and gentamicin in orthopaedic patients was associated with a significant increased rate of post-operative AKI.<sup>7-10</sup> Most of these were uncontrolled before and after studies, but one was a robust quasi-experimental analysis that found the same

effect.<sup>7</sup> In June 2012 within NHS Tayside, policy was changed from flucloxacillin and gentamicin to co-amoxiclav.

This study aimed to assess whether the change in orthopaedic antibiotic prophylaxis policy from flucloxacillin and gentamicin to co-amoxiclav was associated with changes in the rates of post-operative AKI and CDI in adult patients undergoing orthopaedic surgery. The study also aimed to assess the effect of post-operative AKI on length of hospital stay and one year post-operative mortality.

## **Ethical Approval**

Anonymised record linkage was conducted according to the Health Informatics Centre (HIC) standard operating procedure. The Tayside Research Ethics Committee does not require submission of individual studies that follow this standard operating procedure.

## **Methods**

This study included all adults aged  $\geq 18$  years who resided, or died, in the NHS Tayside region and, who underwent an orthopaedic surgical procedure requiring antibiotic prophylaxis, between 1<sup>st</sup> October 2008, and 31<sup>st</sup> December 2013. Patients were identified by the Office of Population, Censuses and Surveys Classification of Surgical Operations and Procedures 4<sup>th</sup> revision (OPCS-4) codes<sup>26</sup> from hospital admissions data (Supplementary Material, Table 1).

Table 1 shows the recommended antibiotics and doses in NHS Tayside before and after the 2012 policy change. Patients who underwent repair of a neck of femur (NOF) fracture received co-amoxiclav as antibiotic prophylaxis throughout the period of observation described here because of concerns raised by orthopaedic surgeons with regard to administering gentamicin in this particular patient group.

[Type here]

This study period encompassed 63 months between October 2008 and December 2013. The study design was an interrupted time series (ITS) with segmented regression analysis<sup>27</sup> using 44 monthly time points before and 19 monthly time points after the intervention in June 2012.

## Definitions

The definition of postoperative renal impairment used the Kidney Disease Improving Global Outcomes (KDIGO) criteria.<sup>28</sup> (Supplementary Material Table 2) Patients with postoperative AKI were classified according to their most severe degree of AKI. This was applied using baseline serum creatinine as the pre-measurement (most recent in the year prior to surgery) and maximal serum creatinine during the first 7 post-operative days as the post-measurement.

The definition of post-operative CDI was any case with a *Clostridium difficile* toxin positive stool sample and/or polymerase chain reaction (PCR) for *Clostridium difficile* positive result within 12 weeks post-operatively. Cases were excluded if there had been another positive sample for *Clostridium difficile* toxin or PCR in the 12 weeks prior to this, due to the possibility of this being a recurrence of CDI.

## Data Sources

Data were provided by HIC<sup>29</sup> at the University of Dundee. Data were linked between the following datasets: Community Health Index (CHI) register, Scottish Morbidity Record of hospital admissions (SMR01), OPCS-coded procedures, laboratory results database, medicines dispensed by community pharmacies, General Register Office death database and Scottish Care Information – Diabetes Collaboration (SCI-DC).

Age, to the nearest year in the year of surgery, sex and social deprivation, the Scottish Index of Multiple Deprivation (SIMD) which is linked to residential postal code, were obtained from the CHI register. A Charlson Comorbidity Index (CCI)<sup>30</sup> score was calculated for each

[Type here]

patient from hospital discharge codes and the number of medication types dispensed from community pharmacies in the previous three months was applied as an additional measure of comorbidity.

Exposure to medicines that predispose to renal impairment (NSAIDs, cyclooxygenase-2 inhibitors, angiotensin-converting enzyme inhibitors (ACE I), angiotensin II receptor antagonists (ARB), diuretics,  $\beta$ -blockers, metformin, other hypoglycaemics and lipid regulating medications), in the three months prior to the operation was ascertained from dispensed prescribing data.

Baseline renal function was obtained from the laboratory database. This was the most recent preoperative serum creatinine measurement in 12 months prior to and including the date of operation, which could include preoperative samples taken during the current admission. The median was calculated if there was more than one serum creatinine measurement taken on the same day. Only patients with both preoperative and postoperative creatinine measurements could be included. The completeness of creatinine data in each group was measured and reported (Table 2). Patients already established on renal replacement therapy pre-operatively were excluded from the analysis. In cases where a patient had a subsequent operation within 7 days of another operation, the second operation was excluded from the analysis.

## **Statistical Analysis**

Monthly rates of AKI were defined by the number of patients with AKI as a proportion of all patients aged  $\geq 18$  years undergoing orthopaedic surgery in each month. Rates were plotted over time for descriptive purposes and the functional form of the relationship before and after the intervention was assessed for linearity with splines. Kolmogorov - Smirnov test<sup>31</sup> and the Shapiro – Wilk test<sup>32</sup> were used to check data were normally distributed. Segmented regression analysis of ITS data was used to assess intervention effects on total AKI at 1, 6, 12

[Type here]

and 18 months after the policy change, with 95% confidence intervals (95%CI)<sup>27, 31</sup> and models were assessed for first order autocorrelation using the Durbin–Watson statistic,<sup>33</sup> with adjustment for autocorrelation using lag terms as required. The effect size at each time point was converted to a relative percentage difference between the observed and predicted (if the intervention had not happened) AKI rate.

To assess for the effect of factors other than policy change on total AKI rates multiple linear regression was used to determine the effects of exposure to medicines that predispose to renal disease, age, sex and CCI on all rates of AKI in the time periods before and after policy change.

The monthly rates of CDI were too small to undergo ITS analysis. Due to there being no or few cases in some months, rates of CDI were calculated using incidence rate ratios<sup>34</sup> (IRR) and 95% CIs.

The median length of stay with IQR was calculated from data extracted from SMR01 data.

Death within 1 year of surgery was calculated as a percentage of all patients who had undergone operations within each group, in patients with and without post-operative AKI.

All analyses were carried out in IBM SPSS (version 21) and SAS (version 9.2.1) software. A p-value of <0.05 was considered statistically significant.

The analysis plan included information that addressed the common risks of bias in ITS studies (Supplementary Material, Table 3).

## **Results**

Data were provided for 14,563 potentially eligible orthopaedic operations in the study period of October 1, 2008 to December 31, 2013. Of these, five duplicate cases were excluded.

[Type here]



Cases were split into two groups, Group 1 (n=12707): patients who underwent orthopaedic operations other than NOF repair, as identified by the International Statistical Classification of Diseases and Related Health Problems ICD9/10 main condition codes, and Group 2 (n=1851): patients who underwent a NOF repair operation. Patients were excluded if they were on the renal replacement register (Group 1 n=33 (0.26%), Group 2 n=8 (0.43%)), if the operation was within 7 days of a previous operation (Group 1 n=22 (0.17%), Group 2 n=0) or if creatinine data were missing (Group 1 n=3410 (26.8%), Group 2 n=81 (4.38%)). In total, 11,004 cases were included (Figure 1).

Compared to patients in the other orthopaedic operation group, patients in the NOF group were older, were more likely to be female and had higher CCI scores (Table 1). However, baseline creatinine was no worse in NOF patients compared with the other orthopaedic operation group, median creatinine 69  $\mu\text{mol/L}$  (IQR 53-94) versus 68  $\mu\text{mol/L}$  (IQR 57-80) respectively.

Table 1 includes the percentage of missing creatinine data within each sub-group. The majority of missing creatinine measurements were post-operative. There were fewer missing creatinine data for patients with NOF repair operations. An examination of the characteristics of patients included *versus* patients who were excluded due to missing data showed that the included patients were, on average, older patients with higher CCI comorbidity scores (Supplementary Material, Table 4).

### **Acute Kidney Injury**

The majority of AKI was mild (Stage 1) AKI (Table 2). Overall, AKI was more common in the NOF repair patients (Group 2) and affected a slightly higher proportion overall in the second study period, while Group 1 had a slightly lower proportion of patient with AKI in the second study period (after prophylaxis changed).

[Type here]

### All other orthopaedic surgeries (Group 1)

The monthly proportion of patients with any stage of post-operative AKI in Group 1 fell following the intervention (Figure 2A) and interrupted time series analysis indicated a relative change of -63% (95% CI -77% to -49%) 18 months after the policy change (Table 3).

Multiple linear regression analyses were performed to assess for the effects of sex, age, CCI and the use of other nephrotoxic drugs. In addition to the policy change only ACE I/ARB were significantly associated with an increase in rates of post-operative AKI ( $\beta=0.186$ ; 95% CI, 0.03 to 0.342;  $p=0.02$ ).

### Neck of Femur repair (Group 2)

Over the whole period of observation AKI rates in this group were highest, and they had most variation in the year preceding the change in antibiotic policy for the other group (Figure 2B). From interrupted time series analysis, there were statistically significant reductions in AKI rates in the following year but these were much smaller relative changes than in Group 1 and there was no significant change at 18 months after the policy change -10%, (95% CI -35% to 15%, Table 3).

Multiple linear regression analyses were performed to assess for the effects of sex, age, CCI and the use of other nephrotoxic drugs. Only ACE I/ARB were significantly associated with an increase in rates of post-operative AKI ( $\beta=0.2520$ ; 95% CI, 0.04 to 0.464;  $p=0.02$ ).

## **Other outcomes**

### Death within 1 year of surgery

A higher proportion of patients with AKI died within 1 year of surgery compared with patients without AKI in both the NOF operations group 36.63% (n=89) versus 25.15%

[Type here]

(n=382), respectively (chi-square=14.09 df=1 p<0.001) and the other orthopaedic operations group 10.48% (n=108) versus 5.18% (n=425), respectively (chi-square=47.33 df=1 p<0.001).

#### Length of hospital stay

In the other orthopaedic operations group the median length of hospital stay was 6 days (IQR 11-3) in patients with post-operative AKI compared with 5 days (IQR 9-3) in patients without post-operative AKI. In the NOF repair operation group was 7 days (IQR 11-4) in patients with post-operative AKI compared with 6 days (IQR 10-3) in patients without post-operative AKI.

#### *Clostridium difficile* infection and SSI Rates

There was no change in the rate of CDI before and after the policy change in either group. The CDI IRR (95% CI) for the NOF repair group was 0.76 (0.28 to 2.08) and for other operations was 0.29 (0.09 to 0.96).

Audits of SSI rates were not done consistently for all types of surgery in the study population throughout the study period; however the SSI rates among the procedures audited were low throughout the study period with no apparent change in rate accompanying the policy changes

### **Discussion**

This large observational study of >14,500 patients, found that a change in prophylactic antibiotic policy from flucloxacillin and gentamicin to co-amoxiclav was associated with decreased rates of post-operative AKI in patients undergoing orthopaedic surgeries (excluding NOF repairs) in the NHS Tayside Region of Scotland, with the majority of patients developing mild (Stage 1) AKI. The association with the change in antibiotic policy

is strengthened because there were much smaller reductions in monthly AKI rates among patients who underwent NOF repair, who received co-amoxiclav throughout the study period, and the change at 18 months post policy change was not significant in this group. There was no change in CDI rates associated with the change in antibiotic policy.

We have shown a reduction in all stages of post-operative AKI by -63% (95% CI -77%,-49%) at 18 months post antibiotic policy change statistically significantly associated with that change in policy. There are likely to be other factors also contributing to the change in rates of AKI, supported by the (smaller magnitude and not significant at 18 months) reduction in NOF patients who were not affected by the policy change. Such factors include anaesthetic practice and post-operative care, such as the Enhanced Recovery Programme, which aims to optimise pre- and post-operative care including reducing the physical stress of the operation, promoting effective analgesia and encouraging early mobilisation and oral nutrition. This programme may lead to an increase in rates of post-operative AKI, as patients may be exposed to more nephrotoxic analgesics and less fluid replacement in the post-operative period than before. There has also been ongoing work throughout our institutions, educating healthcare professionals around recognition, prevention and management of AKI.

The causes of AKI are multifactorial and reported incidence varies dependent on the definition used, as well variations in population groups and settings<sup>35</sup>. The incidence of AKI in acute hospitalised patients is 5-7.5%, with 30-40% of cases occurring in the surgical perioperative period.<sup>36, 37</sup> This is an important contributor to post-operative morbidity and mortality<sup>35</sup> and research suggests that up to 30% of AKI is iatrogenic and/or potentially preventable.<sup>38</sup> The KDIGO AKI guidelines state that major surgery and trauma are exposures for non-specific AKI, where there is no specific kidney disease causing AKI.<sup>28</sup> Therefore, it is important that these patients have their serum creatinine and urine output measured

alongside their clinical status. In line with previous recommendations<sup>7</sup> patients receiving surgical antibiotic prophylaxis are generally undergoing major surgery and, as a minimum, should have serum creatinine measured both pre-operatively and at least 24 hours post-operatively.

The unadjusted one year post-operative mortality was higher in patients who had undergone an NOF repair surgery when compared with those having undergone other orthopaedic surgeries. Within each subgroup crude 1 year post-operative mortality was higher among those with post-operative AKI than those without. This is consistent with the previous findings, including this cohort that AKI contributes to higher mortality.<sup>16, 19, 36-38</sup> We did not, however, find any difference in length of hospital stay, when comparing patients with and without AKI, which may be because the majority of AKI were stage 1 and it is possible that AKI was not detected until after the patient was discharged so did not impact on inpatient management. The potential implications of reduced rates of AKI include reduced mortality, readmission to hospital, progression to CKD and the subsequent socio-economic costs, that have previously been identified in the literature.<sup>20</sup>

There was no change in CDI rates in either group, suggesting that this is not related to changes in surgical prophylactic antibiotic policy. Restrictions on the use of therapeutic antibiotics considered high-risk for CDI, which are used for longer durations, have contributed, alongside infection control procedures and a change in circulating strains and virulence of *Clostridium difficile*,<sup>12, 39</sup> to a reduction in rates of CDI.<sup>40</sup>

Effective restrictions on the use of therapeutic antibiotics associated with high risk for CDI have also been shown in the orthopaedic surgery population, without a subsequent effect on rates of deep SSI.<sup>41</sup> Rates of CDI across the UK have fallen. By 2013, in Scotland there had

274 been reductions in CDI of greater than 79% since 2007/08<sup>42</sup> and in 2013/14, England  
275 reported a reduction in CDI of 76% compared with 2007/08.<sup>43</sup>

276 Prophylactic antibiotics in orthopaedic implant surgery reduces the risk of SSI.<sup>3</sup> In an ageing  
277 population, it is likely that the number of orthopaedic operations is going to increase,  
278 exposing more individuals to prophylactic antibiotics. Any prophylactic antibiotic should aim  
279 to reduce SSI rates, providing cover for commonly implicated pathogens and considering  
280 local resistance patterns, while also minimising the associated risks of AKI, CDI, antibiotic  
281 resistance and allergies. SSI, AKI and CDI are interlinked. The management of SSI is  
282 antibiotic therapy. SSI in severe cases can lead to sepsis. Both exposure to certain antibiotics  
283 and sepsis have been identified as risk factors for non-specific AKI.<sup>28</sup> Further antibiotic  
284 exposure would also put patients at an increased risk of CDI,<sup>24</sup> which itself may result in a  
285 patient becoming dehydrated, another susceptibility for non-specific AKI.<sup>28</sup> It is therefore  
286 important to ensure that intended and unintended consequences of changes in policy are  
287 balanced.

288 The strengths of our study are that we used a robust quasi-experimental study design to  
289 evaluate a real-world policy change, addressed risks of bias for ITS studies in our analysis  
290 plan (Supplementary Material, Table 2), defined operations using OPCS-4 operation  
291 procedure codes (Supplementary Material, Table 3), defined NOF cases using ICD9/10  
292 codes, and used the KDIGO definition of AKI.

293 We addressed the issues common to all non-randomised studies, including potential  
294 ascertainment and selection biases. Data were prospectively entered into electronic databases  
295 and were not subject to manipulation or bias. ITS is the strongest quasi-experimental design  
296 to assess intervention effects in non-randomised setting. It allows for control of trends that

exist before the intervention being studied, by using multiple time points before and after the intervention.<sup>44, 45</sup>

A weakness of our study is that we were unable to adjust for potential effects of intraoperative events and other medications prescribed during in-patient stay on AKI rates because these data are not collected electronically. Furthermore, it is possible that the changes in diagnostic criteria and methods could have masked a true decrease in CDI incidence over the study period.

Rates of post-operative AKI were not assessed in 4% of the NOF repair surgery group and 27% of the other orthopaedic operations group because of missing pre-operative and post-operative creatinine data. These patients were younger and had less comorbidity when compared with patients who were included in the study. This could bias the results in either direction of estimating the rates of AKI. However, there were similar percentages of missing creatinine data in the pre-intervention and post-intervention groups.

Previous literature showed that a prophylactic antibiotic regimen of flucloxacillin and gentamicin in orthopaedic patients was associated with a significant increased rate of AKI. From the previous research carried out by our group it is not clear whether gentamicin or flucloxacillin or both antibiotics increased the risk of post-operative AKI, or by what mechanism.<sup>7</sup> However, this study has shown that using co-amoxiclav for antibiotic prophylaxis in orthopaedic surgery is associated with a decreased rate of post-operative AKI compared to flucloxacillin and gentamicin and is not associated with an increased rate of CDI. This study also raises the importance of monitoring the intended and unintended consequences of policy change within healthcare.

## **Acknowledgements**

[Type here]

320 We acknowledge the support of the University of Dundee for the support and funding of the  
321 Academic Foundation Programme during which Heather Walker carried out analyses for this  
322 work.

### 323 **Funding**

324 This work was supported by the University of Dundee, who funded the Academic Foundation  
325 Programme during which Heather Walker carried out analyses for this work.

### 326 **Transparency declaration**

327 None to declare.

328



1. Plowman R. The socioeconomic burden of hospital acquired infection. *Euro surveillance : bulletin European sur les maladies transmissibles = European communicable disease bulletin* 2000; **5**: 49-50.
2. Whitehouse JD, Friedman ND, Kirkland KB et al. The impact of surgical-site infections following orthopedic surgery at a community hospital and a university hospital: adverse quality of life, excess length of stay, and extra cost. *Infection control and hospital epidemiology* 2002; **23**: 183-9.
3. AlBuhairan B, Hind D, Hutchinson A. Antibiotic prophylaxis for wound infections in total joint arthroplasty: a systematic review. *The Journal of bone and joint surgery British volume* 2008; **90**: 915-9.
4. Scottish Intercollegiate Guidelines Network (SIGN). Antibiotic prophylaxis in surgery. Edinburgh: SIGN, 2014.
5. Uckay I, Hoffmeyer P, Lew D et al. Prevention of surgical site infections in orthopaedic surgery and bone trauma: state-of-the-art update. *The Journal of hospital infection* 2013; **84**: 5-12.
6. Goossens H, Ferech M, Vander Stichele R et al. Outpatient antibiotic use in Europe and association with resistance: a cross-national database study. *Lancet* 2005; **365**: 579-87.
7. Bell S, Davey P, Nathwani D et al. Risk of AKI with gentamicin as surgical prophylaxis. *Journal of the American Society of Nephrology : JASN* 2014; **25**: 2625-32.
8. Challagundla SR, Knox D, Hawkins A et al. Renal impairment after high-dose flucloxacillin and single-dose gentamicin prophylaxis in patients undergoing elective hip and knee replacement. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 2013; **28**: 612-9.
9. Bailey O, Torkington MS, Anthony I et al. Antibiotic-related acute kidney injury in patients undergoing elective joint replacement. *The bone & joint journal* 2014; **96-B**: 395-8.
10. Craxford S, Bayley E, Needoff M. Antibiotic-associated complications following lower limb arthroplasty: a comparison of two prophylactic regimes. *European journal of orthopaedic surgery & traumatology : orthopedie traumatologie* 2014; **24**: 539-43.
11. Bartlett JG. Historical perspectives on studies of *Clostridium difficile* and *C. difficile* infection. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2008; **46 Suppl 1**: S4-11.
12. Carignan A, Allard C, Pepin J et al. Risk of *Clostridium difficile* infection after perioperative antibacterial prophylaxis before and during an outbreak of infection due to a hypervirulent strain. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2008; **46**: 1838-43.
13. Levy EM, Viscoli CM, Horwitz RI. The effect of acute renal failure on mortality. A cohort analysis. *Jama* 1996; **275**: 1489-94.
14. Uchino S, Bellomo R, Goldsmith D et al. An assessment of the RIFLE criteria for acute renal failure in hospitalized patients. *Critical care medicine* 2006; **34**: 1913-7.
15. Chertow GM, Levy EM, Hammermeister KE et al. Independent association between acute renal failure and mortality following cardiac surgery. *The American journal of medicine* 1998; **104**: 343-8.
16. Bell S, Dekker FW, Vadiveloo T et al. Risk of post-operative AKI in patients undergoing orthopaedic surgery- development and validation of a risk score and effect of AKI on survival; an observational cohort study. *BMJ* 2015; **351**.
17. Shusterman N, Strom BL, Murray TG et al. Risk factors and outcome of hospital-acquired acute renal failure. Clinical epidemiologic study. *The American journal of medicine* 1987; **83**: 65-71.
18. Kheterpal S, Tremper KK, Englesbe MJ et al. Predictors of postoperative acute renal failure after noncardiac surgery in patients with previously normal renal function. *Anesthesiology* 2007; **107**: 892-902.

19. Chertow GM, Burdick E, Honour M et al. Acute kidney injury, mortality, length of stay, and costs in hospitalized patients. *Journal of the American Society of Nephrology : JASN* 2005; **16**: 3365-70.
20. Coca SG, Yusuf B, Shlipak MG et al. Long-term risk of mortality and other adverse outcomes after acute kidney injury: a systematic review and meta-analysis. *American journal of kidney diseases : the official journal of the National Kidney Foundation* 2009; **53**: 961-73.
21. Pepin J, Valiquette L, Alary ME et al. Clostridium difficile-associated diarrhea in a region of Quebec from 1991 to 2003: a changing pattern of disease severity. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne* 2004; **171**: 466-72.
22. Health Protection Scotland. Annual report on the surveillance of Clostridium difficile-associated disease (CDAD) in Scotland, October 2007 – September 2008. <http://www.hps.scot.nhs.uk/ewr/article.aspx> (10 March 2015).
23. Public Health England. Clostridium difficile: guidance, data and analysis. <https://www.gov.uk/government/collections/clostridium-difficile-guidance-data-and-analysis#data-submission> (10 March 2015).
24. Hensgens MP, Goorhuis A, Dekkers OM et al. Time interval of increased risk for Clostridium difficile infection after exposure to antibiotics. *The Journal of antimicrobial chemotherapy* 2012; **67**: 742-8.
25. Scottish Antimicrobial Prescribing Group. Guidance to Optimise Antimicrobial Use and Reduce Clostridium difficile Associated Disease in Scottish Hospitals. In: Scottish Government Health Department, ed. Edinburgh, UK, 2008.
26. Health and Social Care Information Centre. Background to OPCS-4 development. <http://systems.hscic.gov.uk/data/clinicalcoding/codingstandards/opcs4/background> (March 26 2015, date last accessed).
27. Wagner AK, Soumerai SB, Zhang F et al. Segmented regression analysis of interrupted time series studies in medication use research. *Journal of clinical pharmacy and therapeutics* 2002; **27**: 299-309.
28. KDIGO Clinical Practice Guideline for Acute Kidney Injury Work Group. *KDIGO Clinical Practice Guideline for Acute Kidney Injury*, 2012.
29. University of Dundee: Health Informatics Centre. <http://www.medicine.dundee.ac.uk/hic> (March 10, 2015 2015, date last accessed).
30. Charlson ME, Pompei P, Ales KL et al. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *Journal of chronic diseases* 1987; **40**: 373-83.
31. Zhang F, Wagner AK, Soumerai SB et al. Methods for estimating confidence intervals in interrupted time series analyses of health interventions. *Journal of Clinical Epidemiology* 2007; **62**: 143-8.
32. Shapiro SS, Wilk MB. An analysis of variance test for normality (complete samples). *Biometrika* 1965; **52**: 3-4.
33. Durbin J, Watson GS. Testing for serial correlation in least squares regression. I. *Biometrika* 1950; **37**: 409-28.
34. Rotham KJ GS. *Modern edipdemiology*. Philadelphia:: Lippincott Williams & Wilkins, 2008.
35. Davenport A, Stevens P. UK Renal Association Clinical Practice Guidelines: Module 5 Acute Kidney Injury. 2008.
36. Thakar CV, Christianson A, Freyberg R et al. Incidence and outcomes of acute kidney injury in intensive care units: a Veterans Administration study. *Critical care medicine* 2009; **37**: 2552-8.
37. Uchino S, Kellum JA, Bellomo R et al. Acute renal failure in critically ill patients: a multinational, multicenter study. *Jama* 2005; **294**: 813-8.
38. Stevens PE, Tamimi NA, Al-Hasani MK et al. Non-specialist management of acute renal failure. *QJM : monthly journal of the Association of Physicians* 2001; **94**: 533-40.
39. National Institute for Health and Care Excellence (NICE). Clostridium difficile infection: risk with broad-spectrum antibiotics. National Institue for Health and Care Excellence, 2015; 30.
40. Vernaz N, Hill K, Leggeat S et al. Temporal effects of antibiotic use and Clostridium difficile infections. *The Journal of antimicrobial chemotherapy* 2009; **63**: 1272-5.

41. Al-Obaydi W, Smith CD, Foguet P. Changing prophylactic antibiotic protocol for reducing Clostridium difficile-associated diarrhoeal infections. *Journal of orthopaedic surgery* 2010; **18**: 320-3.
42. Connaghan J. Update to the Clostridium difficile HEAT target. In: Scottish Government General Health & Social Care Department, ed, 2013.
43. Annual Epidemiological Commentary : Mandatory MRSA, MSSA and E. coli bacteraemia and C. Difficile infection. In: Department of Health, ed, 2014.
44. Higgins JPT RC, Reeves BC, Deeks JJ, Shea B, Valentine JC, Tugwell P, Wells G. . Issues relating to study design and risk of bias when including non-randomised studies in systematic reviews on the effects of interventions. *Res Synth Methods* 2012: 12–25.
45. Ramsay CR, Matowe L, Grilli R et al. Interrupted time series designs in health technology assessment: lessons from two systematic reviews of behavior change strategies. *International journal of technology assessment in health care* 2003; **19**: 613-23.

Table 1. Baseline characteristics and missing creatinine data

Characteristic	Operations other than NOF repair		NOF repair operations	
	Before 2012 policy change	After 2012 policy change	Before 2012 policy change	After 2012 policy change
<b>Recommended antibiotics (dose)</b>	Flucloxacillin (1g) x 4 doses + one dose Gentamicin (4mg/kg)	3 doses Co-amoxiclav (1.2g)	3 doses Co-amoxiclav (1.2g)	3 doses Co-amoxiclav (1.2g)
<b>Patients (n)</b>	6267	2975	1250	512
<b>Mean Age (yr) (SD)</b>	68.5 (14)	68.9 (13.6)	80 (11.4)	80.6 (10.2)
<b>Median Baseline SCr (<math>\mu\text{mol/L}</math>) (IQR)</b>	74 (62-88)	68 (57-80)	72 (57-94)	69 (53-94)
<b>Sex n(%)</b>				
<b>Female</b>	3619 (57.7)	1774 (59.6)	914 (73.1)	372 (72.7)
<b>Male</b>	2648 (42.3)	1201 (40.4)	336 (26.9)	140 (27.3)
<b>SIMD n(%)</b>				
<b>1-3 (most deprived)</b>	1212 (19.3)	573 (19.3)	244 (19.5)	113 (22.1)
<b>4-7</b>	2679 (42.7)	1299 (43.7)	545 (43.6)	217 (42.4)

[Type here]

<b>8-10 (least deprived)</b>	2269 (36.2)	1044 (35.1)	444 (35.5)	171 (33.4)
<b>CCI n(%)</b>				
<b>Low (0)</b>	5282 (84.3)	2317 (77.9)	789 (63.1)	303 (59.2)
<b>Medium (1 or 2)</b>	870 (13.9)	584 (19.6)	400 (32)	183 (35.7)
<b>High (≥3)</b>	115 (1.8)	74 (2.5)	61 (4.9)	26 (5.1)
<b>Patients before missing creatinine exclusions n(%)*</b>	8515	4137	1304	539
<b>Missing pre-operative creatinine</b>	453 (5.32)	192 (4.64)	7 (0.54)	4 (0.74)
<b>Missing post-operative creatinine</b>	1995 (23.43)	1039 (25.11)	49 (3.76)	22 (4.08)
<b>Missing both pre and post-operative creatinine</b>	405 (4.76)	173 (4.18)	4 (0.31)	2 (0.37)

\* Data are presented as n (%) of all patients after exclusion of duplicates, patients on RRT and subsequent operation within 7 days

[Type here]

Table 2. AKI rates by each stage


AKI Stage n(%)	Operations other than NOF repair		NOF repair operations	
	Before 2012 policy change	After 2012 policy change	Before 2012 policy change	After 2012 policy change
<b>1</b>	618 (9.86)	239 (8.03)	153 (12.24)	67 (13.09)
<b>2</b>	95 (1.52)	22 (0.74)	16 (1.28)	4 (0.78)
<b>3</b>	45 (0.72)	12 (0.4)	2 (0.16)	1 (0.2)
<b>None</b>	5509 (87.9)	2702 (90.8)	1079 (86.32)	440 (85.94)

[Type here]

**Table 3. ITS analyses intervention effects at 1, 6, 12 and 18 months post-policy change for other orthopaedic operations and NOF repair operations, all AKI Stages (Rate per 100 patients)**

	Operations other than NOF repair				NOF repair operations			
	1 month	6 months	12 months	18 months	1 month	6 months	12 months	18 months
<b>Absolute difference</b>	-6.1	-8.2	-10.6	-13.1	-4.7	-3.8	-2.8	-1.8
<b>95% CI</b>	(-9.7,-2.6)	(-11.2,-5.2)	(-13.9,-7.4)	(-17.3,-8.9)	(-8.8,-0.5)	(-7.3,-0.4)	(-6.5,0.8)	(-6.6,3.0)
<b>Relative change</b>	-36.0%	-45.2%	-54.8%	-63.2%	-28.7%	-22.8%	-16.1%	-10.0%
<b>95% CI</b>	(-54.6%, -17.4%)	(-57.7%, -32.6%)	(-65.3%, -44.3%)	(-77.3%, -49.1%)	(-52.2%, -5.3%)	(-40.8%, -4.8%)	(-34.6%, 2.3%)	(-34.7%, 14.8%)


Figure 1: Summary of case groupings and exclusions



**Figure 2A: AKI (Stages 1-3 combined) in orthopaedic operations other than NOF repair from October 2008 to December 2013. The Policy Change was from flucloxacillin plus gentamicin to co-amoxiclav for antibiotic prophylaxis.**

[Type here]





**Figure 2B: AKI (Stages1-3 combined) in NOF repair operations from October 2008 to December 2013 with co-amoxiclav the recommended antibiotic prophylaxis throughout. The Policy Change was for operations other than NOF repair.**